

## Introduction

Necrotizing fasciitis (NF) is a rare, complex disease with a mortality rate between 25-35%. Typically the infecting agent invades the soft tissues, aggressively attacking deep soft tissue layers with resultant liquefaction within hours of the initial exposure. Early diagnosis is key in lowering mortality rates. However, disease rarity, along with the inverse relationship of physical exam to symptom severity may contribute to a delay in diagnosis and treatment. The gold standard treatment includes emergent fasciectomy, serial débridements, broad spectrum antibiotics, local wound care, and eventual skin grafting.

In this case study, a 49-year-old male presented with acute NF of the left lower extremity. A novel antimicrobial formula, Hexagen gel, was employed in the treatment protocol.

## Laboratory Risk Indicator for Necrotizing Fasciitis (LRINEC)

| VARIABLE                                      | UNITS                       | SCORE               |
|---|-----------------------------|---------------------|
| C-reactive protein                            | ≥150 mg/L                   | 4 points            |
| White blood cell count (per mm <sup>3</sup> ) | 15-25<br>>25                | 1 point<br>2 points |
| Hemoglobin                                    | 11.0-13.5g/dl<br>< 11 g/dl  | 1 point<br>2 points |
| Serum sodium                                  | ≥135 mmol/l<br><135 mmol/l  | 1 point<br>2 points |
| Serum creatinine                              | >1.6 mg/dl (or >141 μmol/l) | 2 points            |
| Serum glucose                                 | >180 mg/dl (or > 10 mmol/L) | 1 point             |

\*A LRINEC score of ≥ 8 correlates with a 93.4% likelihood of NF infection.

## Case Report

### 12/9/16 (7:29pm)

- 49-year-old Hispanic male with complaints of shortness of breath and left leg pain.
- Patient was hypotensive, tachycardic, hyponatremic, and hyperglycemic; a small wound was present on his anterior left leg due to a splinter two days prior.
- White blood cell count (WBC) was  $23 \times 10^3$  cells/mL, serum creatinine was elevated, and hemoglobin decreased, giving a LRINEC score of 8.
- Diagnosis: Septic shock secondary to NF of the left lower extremity.
- Treatment: To OR for emergent débridement and fasciectomy.

### 12/9/16 (11:00pm)

- Intra-op: Erythema, edema, and liquefaction of LLE.
- Extensive incisions from the mid thigh medially and laterally down to the ankle. Significant purulence in the subcutaneous tissues and surrounding fascia.
- Post-op: WBC rose to  $25 \times 10^3$  cells/mL. Wound cultures positive for beta-hemolytic group A strep. Broad spectrum IV antibiotics begun.

### 12/12/17

- Follow-up débridement and salvage procedure with Versajet. Negative-pressure therapy applied to avoid fluid collection. Post-op IV antibiotics continued.
- WBC:  $30 \times 10^3$  cells/mL. Concern about viability of limb.

### 12/16/17

- Further débridement. Wound dressed with Hexagen antimicrobial gel, Adaptic, and Kerlix.
- Post-surgical WBC reduced to  $22 \times 10^3$  cells/mL.
- Post-op IV antibiotics continued. Daily dressing changes with Hexagen, Adaptic, and Kerlix.
- Wound improved; WBC decreasing.

### 12/28/17

- More intra-op débridement performed with application of split-thickness skin grafts harvested from right thigh. Hexagen, Adaptic, and negative-pressure therapy applied to the grafted wound base.
- IV antibiotics continued. Post-surgical WBC reduced to normal ( $10 \times 10^3$  cells/mL).

### 1/2/17

- Patient discharged from hospital with daily dressing changes of Hexagen, Adaptic, Kerlix, and light ACE wraps.

### 2/16/17

- 6-week post-op evaluation: Patient returned to work as a construction worker at a normal activity level.
- Patient continues to apply Hexagen and Kerlix daily; happy with his post-op course.

## Results



## Results cont.



## Discussion

Necrotizing fasciitis (NF) is a limb and life-threatening soft-tissue infection with acute onset and rapid progression. The rapidly progressive nature of NF requires prompt and aggressive surgical débridement, tissue reconstruction or amputation may be required. The elevated WBC on admission decreased rapidly following the application of Hexagen antimicrobial gel. Hexagen antimicrobial gel may serve as a useful tool for ancillary antimicrobial therapy in NF infections.

## Conclusion

Further study is needed to justify use of Hexagen antimicrobial therapy for NF. Our case report suggests that its use as ancillary therapy yields better outcomes than traditional treatment methods alone.

## References

- Wong CH, Wang YS. The diagnosis of necrotizing fasciitis. *Curr Opin Infect Dis* 2005;18:101-6.
- Anaya DA, McMahon K, Nathens AB, Sullivan SR, Foy H, Bulger E. Predictors of mortality and limb loss in necrotizing soft tissue infections. *Arch Surg* (2005) 140:151-7. doi:10.1001/archsurg.140.2.151
- Wong, C., Khin, L., Heng, K., Tan, K., & Low, C. (2004). The LRINEC (Laboratory Risk Indicator for Necrotizing Fasciitis) score: A tool for distinguishing necrotizing fasciitis from other soft tissue infections\*. *Critical Care Medicine*, 32(7), 1535-1541. doi:10.1097/01.ccm.0000129486.35458.7d
- Bilton BD, Zibari GB, McMillan RW, Aultman DF, Dunn G, McDonald JC. Aggressive surgical management of necrotizing fasciitis serves to decrease mortality: a retrospective study. *Am Surg* 1998;64:397-400.
- Wong CH, Yam AK, Tan AB, Song C. Approach to debridement in necrotizing fasciitis. *Am J Surg* (2008) 196:19-24. doi:10.1016/j.amjsurg.2007.08.076
- Menichetti F, Giuliano S, Fortunato S. Are there any reasons to change our behavior in necrotizing fasciitis with the advent of new antibiotics? *Curr Opin Infect Dis* (2017), 30:172-179 DOI:10.1097